

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Fig. 1A

Survival distributions, YF-VAX® and ChimeriVax™-JE construct with and without a mutation at E279 (Met→Lys). 4 day-old suckling mice. The dose is approximately equivalent across viruses tested.

A. Dose ~0.7 log₁₀ PFU




Log rank test $p < 0.0001$, including YF-VAX

Log rank test $p = 0.1216$ comparing the three ChimeriVax™-JE constructs

Fig. 1B


Dose ~1.7 log₁₀ PFU



Log rank test p= 0.0016

Fig. 1C

Dose ~2.7 log₁₀ PFU



Log rank test p= 0.0558

Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7

